

Neutrinos: Summarizing the State-of-the-Art

J. W. F. Valle

Instituto de Física Corpuscular – C.S.I.C., Universitat de València
Edificio Institutos, Aptdo. 22085, E-46071 València, Spain
E-mail:valle@ific.uv.es

Abstract. I review oscillation solutions to the neutrino anomalies and discuss how to account for the required pattern of neutrino masses and mixings from first principles. Unification and low-energy bottom-up approaches are discussed, the latter open up the possibility of testing neutrino mixing at high energy colliders, such as the LHC. Large ν_e mixing is consistent with Supernova (SN) astrophysics and may serve to probe galactic SN parameters at Cherenkov detectors. I discuss the robustness of the atmospheric neutrino oscillation hypothesis against the presence of Flavor Changing (FC) Non-Standard neutrino Interactions (NSI), generally expected in models of neutrino mass. Atmospheric data strongly constrain FC-NSI in the ν_μ - ν_τ channel, while solar data can be explained by FC-NSI in the ν_e - ν_τ channel, or, alternatively, by spin flavor precession. I illustrate how a neutrino factory offers a unique way to probe for FC-NSI and argue that a near-site detector is necessary in order to probe for leptonic mixing and CP violation.

1 Solar Neutrinos [1,2,3,4,5,6]

For decades, the number of solar neutrinos detected in underground experiments [1] has been less than expected from theories of energy generation in the sun [2], suggesting that either the understanding of the Sun was incomplete, or that the neutrinos were changing from one type to another in transit from the core of the Sun to the detector. At the moment we have a conclusive proof for the latter possibility, though we still can not pin down the exact neutrino conversion mechanism implied.

Solar neutrinos have now been detected through the geochemical method (the $\nu_e + {}^{37}\text{Cl} \rightarrow {}^{37}\text{Ar} + e^-$ reaction at the Homestake experiment and the $\nu_e + {}^{71}\text{Ga} \rightarrow {}^{71}\text{Ge} + e^-$ reaction at the Gallex, Sage and GNO experiments) and also directly using Cherenkov techniques at Kamiokande and SuperKamiokande (SK) and the Sudbury Neutrino Observatory (SNO). At SK the reaction is $\nu_e e$ scattering on water, while SNO detects neutrinos using 1000 tonnes of heavy water (D₂O) and is sensitive to the charged current (CC) and neutral current (NC), as well as to the elastic scattering (ES) reaction.

At present all experiments observe a deficit of 30 to 60 % with an energy dependence mainly due to the lower Chlorine rate. The high statistics of SK after 1496 days of data-taking also provides very useful information on the recoil electron energy spectrum and this is presently well described by the flat hypothesis, placing important restrictions on neutrino parameters. Moreover, SK measures the zenith angle distribution (day/night effect) which is sensitive to the effect

of the Earth matter in the neutrino propagation, observing no significant effect. The absence of a clear hint of spectral distortion, day-night or seasonal variation implies that, to a large extent the solar neutrino problem rests heavily on the rate discrepancy and, from this point of view, future experiments are most welcome.

The global status of the interpretation of the solar neutrino data in terms of active neutrino oscillations is illustrated in Fig. 1. This figure refers to the 1258




Fig. 1. The solar χ^2 profiles versus angle and Δm^2 , from [3], before [5,6].

day SK data sample. As can be seen the best of the oscillation solutions is the Large Mixing Angle (LMA) solution, confirming first pre-SNO hints obtained in ref. [4]. This was derived solely on the basis of the observed flatness of the recoil electron energy spectrum of SK, highlighting its importance in shaping up what is now much more strongly supported.

Solar neutrinos from the decay of ^8B have been detected at SNO via the CC reaction on deuterium and by the ES reaction. Such CC reaction is sensitive only to ν_e 's, while the ES reaction also has a small sensitivity to ν_μ 's and ν_τ 's. SNO has recently observed also NC neutrino interactions [5].

Using the standard ^8B shape they measure the ν_e component of the ^8B solar flux ϕ_e above 5 MeV as well as the non- ν_e component $\phi_{\mu\tau}$. As shown in Fig. 2 the latter is significantly above zero, providing strong evidence for solar ν_e conversion into another active neutrino flavor. This shows the flux of ^8B solar neutrinos which are μ or τ flavor versus the ν_e flux derived from the three neutrino reactions in SNO. The diagonal bands show the total ^8B flux as predicted by the SSM [2] (dashed lines) and that measured with the NC reaction in SNO (solid band). The intercepts of these bands with the axes represent the $\pm 1\sigma$ errors. The bands intersect at the fit values for ϕ_e and $\phi_{\mu\tau}$, indicating that the combined flux results are consistent with neutrino flavor conversion with no distortion in the ^8B neutrino energy spectrum. They also find the total flux measured with the NC reaction to be consistent with solar models.

SNO has also measured day and night solar neutrino energy spectra and rates [6]. They give an analysis in terms of matter-enhanced oscillations of two active flavors showing how it strongly favors the LMA solution. The update of global fits including these latest data worsen the status of non-LMA and especially that




Fig. 2. SNO neutral current measurement [5].

of non-active solutions, see ref. [2,7]. We eagerly await a possible confirmation of the LMA hypothesis at KamLAND [8]. As we will see later there are also good active non-oscillation neutrino conversion mechanisms based on flavor-changing neutrino NSI or spin flavor precession that can account well for the data.

2 Atmospheric Neutrinos [9,10,11,12,13,14]

Atmospheric neutrinos are produced as the decay products in hadronic showers from particles produced in collisions of cosmic rays with air in the upper atmosphere. These have been observed in several experiments [9]. Although individual ν_μ or ν_e fluxes are only known to within 30% accuracy, their ratio is predicted to within 5% over energies varying from 0.1 GeV to 100 GeV [10]. There has been a long-standing discrepancy between the predicted and measured μ/e ratio of the muon ($\nu_\mu + \bar{\nu}_\mu$) over the electron atmospheric neutrino flux ($\nu_e + \bar{\nu}_e$) both in water Cherenkov experiments (Kamiokande, Super-Kamiokande and IMB) as well as in the iron calorimeter Soudan2 experiment. Moreover a strong **zenith-angle dependence** has been observed both in the sub-GeV and multi-GeV energy range, but only for μ -like events, the zenith-angle distributions for the e -like being consistent with expectation. Such clear deficit of neutrinos coming from below is very suggestive of ν_μ oscillations. Zenith-angle distributions have also been recorded for upward-going muon events in SK and MACRO which are also consistent with the ν_μ oscillation hypothesis.

Of all laboratory searches for neutrino oscillation, reactors provide the highest sensitivity in Δm_{32}^2 and thus are most relevant to confront with the data from underground experiments. The restrictions on Δm_{32}^2 and $\sin^2(2\theta_{13})$ that follow from the non observation of oscillations at the Chooz reactor experiment are shown in Fig. 3. The curves represent the 90, 95 and 99% CL excluded region defined with 2 d. o. f. for comparison with the Chooz published results. For large Δm_{32}^2 this gives a stringent limit on $\sin^2(2\theta_{13})$, but not for low Δm_{32}^2 values. After combining with the full body of atmospheric neutrino data, one finds a stringent bound $\sin^2(\theta_{13}) < 0.045$ at 99% CL [12]. Neglecting θ_{13} we display in Fig. 4




Fig. 3. Region in Δm_{32}^2 and $\sin^2(2\theta_{13})$ excluded by the Chooz reactor, from [11].

the allowed regions of atmospheric oscillation parameters Δm_{32}^2 – $\sin^2(2\theta_{23})$ obtained in a global analysis. For previous analyzes, see [13] and references therein.




Fig. 4. Atmospheric oscillation parameters from global analysis in [14].

3 Supernova Neutrinos [15,16].

The large-angle neutrino oscillation between the electron neutrino ν_e favored by current solar neutrino data must be properly taken into account in the interpretation of the SN neutrino data. Indeed, it will induce a significant deformation of the energy spectrum of neutrinos coming from a supernova (SN) explosion and on this basis it has been argued [17] to be at odds with the historical observation of neutrinos from SN1987 [18].

The impact of $\bar{\nu}_e \leftrightarrow \bar{\nu}_{\mu,\tau}$ neutrino oscillations on the observed $\bar{\nu}_e$ signal of supernova SN 1987A can be studied by performing a maximum-likelihood analysis. The fit parameters are the released binding energy E_b and the average neutrino energy $\langle E_{\bar{\nu}_e} \rangle$. It was found that $\bar{\nu}_e \leftrightarrow \bar{\nu}_{\mu,\tau}$ oscillations with large mixing angles have lower best-fit values for $\langle E_{\bar{\nu}_e} \rangle$ than small-mixing angle (SMA) oscillations. Moreover, the inferred value of $\langle E_{\bar{\nu}_e} \rangle$ is already lower than found in simulations in the SMA or no-oscillation case. This apparent conflict has been

interpreted as evidence against the large mixing oscillation solutions to the solar neutrino problem. In order to quantify the degree to which the SN data favor no-oscillations over the large mixing solutions their likelihood ratios were used as well as a Kolmogorov-Smirnov test. The result was that within the range of SN parameters predicted by simulations there are regions where the LMA solution is either only marginally disfavored or even favored over the SMA case [15]. This implies that the LMA solution is not in conflict with the current understanding of SN physics. In contrast, vacuum oscillation and LOW solutions may be excluded at the 4σ level for most of the SN parameter ranges found in simulations. The reason for the difference amongst the large mixing solutions has to do with the role of ν_e regeneration in the Earth matter.

One can go a step further, namely to perform a combined likelihood analysis of the observed SN1987A neutrino signal and of the solar neutrino data [3]. In the calculation of neutrino survival and conversion probabilities a power-law $1/r^3$ profile typical of supernova envelopes was used and the level-crossing probability was determined analytically in a form valid for all mixing angles [19]. It has been found that, although the spectrum swap argument is sufficient to rule out vacuum-type solutions for most reasonable choices of astrophysics parameters, the LOW solution may still be acceptable. On the other hand it was found that the LMA solution can easily survive as the best overall solution, although its size is typically reduced when compared to solar-only fits [3]. How about a future supernova?

The significant distortion of the energy spectrum of neutrinos coming from a supernova (SN) explosion expected by active LMA ν_e oscillations can be used to extract some of the properties of SN driven by gravitational collapse. In ref. [16] a simple but powerful method was proposed to extract the original temperatures of both anti-electron and muon/tau neutrinos at the neutrinosphere from the observation of anti-electron neutrinos at a large water Cherenkov detector such as Super-Kamiokande. Fig. 5 illustrates the potential of the method in determining intrinsic SN parameters. As can be seen one obtains a rather good determina-




Fig. 5. LMA ν_e oscillations allow a determination of SN parameters [16].

tion of the important astrophysical parameter $\tau \equiv \langle E_{\bar{\nu}_\mu} \rangle / \langle E_{\bar{\nu}_e} \rangle$ characterizing the non-electron flavor SN neutrino spectra. The result is rather robust in the sense that it is not too sensitive to astrophysical assumptions such as energy equipartition. Prospects are substantially better for the projected megaton-class SK successor, HyperKAMIOKANDE.

4 Basic Neutrino Theory [20,21,22,23,24,25,26,27,28,29]

The basic gauge theoretical aspects of neutrinos have been developed in the early eighties [20,21,22,23,24,25,26,27,28,29]. The other basic step was the formulation of the MSW effect itself [30]. With both ingredients one can describe all the exciting experimental developments of the past few years [31]. In order to make sense of present neutrino data through neutrino oscillations, neutrinos should have mass. In analogy with the Kobayashi-Maskawa (KM) quark mixing matrix [32], let K denote the lepton mixing matrix. The simplest form for K in a gauge theory of the weak interaction is characterized by

- the atmospheric angle $\theta_{23} \equiv \theta_A$
- the solar angle $\theta_{12} \equiv \theta_S$
- the reactor angle $\theta_{13} \equiv \theta_R$
- 1 KM-like CP phase ϕ_{13}
- 2 extra (Majorana) phases ϕ_1, ϕ_2 [23]

The “Majorana” phases are physical [24]. Although they do not show up in ordinary ($\Delta L = 0$) oscillations, they affect $\Delta L = 2$ processes. Unfortunately, however, the effect is tiny due to helicity suppression, reflecting the V-A nature of the weak interaction [24]. Also the effect of the “Dirac” CP violation due to ϕ_{13} is small since it disappears as $\theta_{13} \rightarrow 0$ [33]. A new generation of long-baseline neutrino oscillation experiments using a neutrino beam from the decay of muons in a storage ring is being considered [34]. Studying θ_{13} and the resulting leptonic CP violating effects provides a challenge for such so-called **neutrino factories**.

The structure of charged and neutral current weak interactions is more complex in models of neutrino mass containing $SU(2) \otimes U(1)$ singlet leptons [23]: the CC is characterized by a rectangular (effectively non-unitary) matrix, while the NC is described by a projective matrix with non-diagonal entries coupling different mass eigenstate neutrinos with each other. Similarly, if there are light sterile neutrinos, the nontrivial structure of K must be taken into account.

Current solar and atmospheric data can be understood well with the usual three massive neutrino flavors. The information on the neutrino oscillation parameters may be summarized by saying that θ_{23} is nearly maximal, θ_{12} is large, but non-maximal, and the reactor angle θ_{13} is rather small. In contrast there is at the moment no information on the CP phases. It is clear from the above that the atmospheric mass splitting is much larger than the solar one, $\Delta m_{atm}^2 \gg \Delta m_\odot^2$. Such hierarchical pattern can be realized in two ways, illustrated in Fig. 6. Last, but not least, the oscillation data are sensitive only to neutrino mass splittings, not to the absolute scale of neutrino mass. Nor do they have any bearing on the




Fig. 6. Normal and inverse-hierarchical neutrino masses. The wide and narrow bands represent the atmospheric and solar mass splittings, respectively.

fundamental issue of whether neutrinos are Dirac or Majorana particles. This brings us to the significance of the $\beta\beta_{0\nu}$ decay [35] in deciding the nature of neutrinos. The connection between the two is given by the **black-box theorem** which states that, in a “natural” gauge theory, whatever the mechanism for inducing $\beta\beta_{0\nu}$ is, it is bound to also yield a Majorana neutrino mass at some level, and vice-versa, as illustrated by Fig. 7. Quantifying⁰⁻⁰ the implications of




Fig. 7. The black-box $\beta\beta_{0\nu}$ argument, from [27].

the black-box argument is a model-dependent enterprise, but can be done once a particular model is specified. It serves at least to highlight the importance of confirming or refuting the present hint [36,37] in a more sensitive experiment such as GENIUS [38].

Can one predict neutrino properties from first principles? Such important enterprise is unfortunately difficult, for at least three basic reasons. First the underlying **scale** is unknown, models using anything between the Planck or String scale, through the $E(6)$ or $SO(10)$ GUT scale, or an intermediate scale associated with Peccei-Quinn or Left-Right symmetry, all the way down to the weak $SU(3) \otimes SU(2) \otimes U(1)$ scale itself. Second, the **mechanism** is unknown, and there are both tree level, like the so-called seesaw schemes [21,23,25,26] as well as radiative [39] or hybrid alternatives. Similarly the status of B-L (baryon minus lepton number) symmetry is unknown, it may either be a broken gauge symmetry, such as in the left-right or $SO(10)$ schemes, or simply a global ungauged symmetry, in which case the corresponding Goldstone boson – called majoron – is physical [25,26]. Last, but not least, there is **no theory of flavour**, although many

Yukawa textures have been suggested which would arise from suitable family symmetries.

Renormalizable neutrino masses are forbidden by the Standard Model (SM), given its restricted field content and symmetries. The simplest and most basic way to add neutrino masses is to regard the SM only as the effective low-energy limit of some, unknown, higher energy theory, which may well contain an L-violating dimension-five term [20]

$$\mathcal{L}_5 = \frac{\lambda_{ij}}{M_5} (\ell_i \phi)(\ell_j \phi), \quad (1)$$

involving the lepton doublet fields ℓ_i ($i, j = e, \mu, \tau$), and the Higgs doublet field ϕ . Here M_5 is an arbitrary mass scale, related to the unknown high energy theory




Fig. 8. Neutrino masses from dimension-five operator [20].

of which the SM is an effective theory and λ_{ij} are dimensionless couplings. After electroweak symmetry breaking, eq. (1) leads to neutrino Majorana masses m_ν^{ij} given by

$$m_{ij}^\nu = \frac{\lambda_{ij} v^2}{M_5}. \quad (2)$$

where $v = 174$ GeV is the Higgs vacuum expectation value (vev). Note that eq. (2) explicitly violates lepton number symmetry.

One way to realize this picture is in terms of the so-called seesaw scenario for neutrino mass [21,22,23,25,26], where this operator is induced either through the exchange of heavy $SU(3) \otimes SU(2) \otimes U(1)$ singlet neutrinos or extra scalar bosons related, say, with the breaking of extended left-right-type gauge symmetries. The first challenge for unification seesaw schemes is that the smallness of the quark KM mixing angles typically suggests that leptonic mixing angles should also be small. However this is certainly possible to avoid owing to the lack of predictivity. Indeed, despite its attractiveness and the existence of many papers claiming to have “predictive” seesaw models of neutrino mass and mixing [40], such models at most **post-dict** the observed angles and splittings and it is hardly possible to obtain “smoking-gun” tests of their validity.

5 Neutrino Models

Notwithstanding the limitations of theory in predicting neutrino properties from first principles, I will mention some recent examples of predictive theoretical schemes for accounting for the neutrino anomalies. Given the vast literature [40,41], I must confine here to a personal selection of recent neutrino models.

5.1 Neutrino Mass Unification [42]

The idea of neutrino mass unification is inspired by the paradigm of unification of fundamental interactions. This “minimalistic” ansatz postulates that the neutrino mass and mixings observed at low energies take a very simple form at some high energy **neutrino unification** scale M_X . The model adds to the basic Lagrangian the dimension-five operator in eq. (1) assumed to be characterized at the scale $M_5 = M_X$ by a single real dimension-less parameter λ_0 . In this picture neutrino masses arise from a common seed at M_X , while their splittings are induced by renormalization effects, as illustrated in Fig. 9. This




Fig. 9. Artists’ view of neutrino masses arising from a common seed (1 eV) at a large neutrino unification scale $M_X \sim 10^{14}$ GeV [42].

model is inconsistent with small solar mixing, thus it “predicts” a large, though non-maximal solar angle, according to the formula

$$\Delta m_S^2 \approx 2\Delta m_A^2 [\cos 2\theta_S + 2\sin^2\theta_R],$$

where Δm_S^2 and Δm_A^2 are the solar and atmospheric splittings while θ_S and θ_R are the solar and reactor angles, respectively. Neutrinos can lie in the eV range and thus be seen in tritium beta decays [43] as well as contribute to the hot dark matter [44]. In contrast to other mechanisms leading to quasi-degenerate neutrinos [45,46] which may be unnatural [47], here this is not the case because opposite neutrino CP signs [29,48,49] suppress the rate for neutrinoless double beta decay. Taken at face value, an analysis including the most recent NC SNO data [2] suggests that the preferred solution in this case will be in terms of vacuum oscillations, instead of LMA, which as we saw is disfavored by SN1987A. In order to evaluate more carefully our prediction a three-neutrino global fit of all neutrino data, similar to that in [11], would be required.

5.2 Minimalistic Seesaw Plus Gravity [50]

The idea here rests on the observation that gravity alone can account for the solar neutrino data by Planck-mass L-violating effects, while the atmospheric neutrino anomaly can be due to the existence of a single right-handed neutrino, instead of three [33]. One assumes it to be at an intermediate mass scale between 10^9 GeV and 10^{14} GeV. To first approximation only one of the three flavor neutrinos acquires mass *a la seesaw*, identified as the atmospheric scale. The other two neutrinos remain massless, as illustrated in Fig. 10 (left panel). Their degeneracy is lifted only by tiny gravitational effects (right panel) due to the dimension-five operator. Such simple scheme solves the current solar and




Fig. 10. Minimalistic neutrino masses. The degeneracy of the massless states on the left is lifted by gravitational effects, illustrated in the right panel.

atmospheric neutrino puzzles with probably the smallest amount of beyond-the-Standard-Model ingredients. Note that although the neutrino mixing angles are not exactly predicted, they can be large, which agrees well with the current experimental situation. However it requires the vacuum or just-so solution to the solar neutrino problem, not currently favored. One test for the model is the search for anomalous seasonal effects at Borexino [51]₀₋₀.

5.3 Supersymmetry As the Origin of Neutrino Mass [52,53]

The suggestion that neutrino masses may be genuinely supersymmetric is rather old [54,55]. One simply postulates that they arise as a result of the violation of R parity (RPV),⁰⁻⁰

$$R_p = (-1)^{3B+L+2S}$$

where S, B, L denote spin, baryon and total lepton number, respectively. Standard model particles (including Higgs scalars) are RP-even, while their superpartners are odd. R parity acts as a selection rule, according to which supersymmetric particles can only be pair-produced, the lightest **sparticle** being stable. These properties have been taken to be the basis of most searches of supersymmetric particles in the laboratory. However, R parity is by no means a sacred symmetry and its violation could substantially affect the expected SUSY signatures while endowing neutrinos with mass.

There are several ways to realize this as illustrated in Fig. 11. The violation




Fig. 11. Ways to break R parity.

of R-parity can arise explicitly [55] as a residual effect from unification that filters down to low energies, or simply as an electroweak phenomenon generated due to non-zero $SU(2) \otimes U(1)$ singlet sneutrino vevs [56]. The latter may be realized either with gauged or ungauged lepton number, in the latter case there is a majoron.

Barring the existence of special symmetries, when R-parity breaks explicitly there can be both bi-linear and tri-linear terms in the low-energy superpotential

$$W = W_{MSSM} + \mu_\alpha \ell_\alpha H_u + \text{trilinears}.$$

However there may exist symmetries which simplify the otherwise very complex structure of explicit RPV terms at low-energy, characterized by too many new independent coupling constants. An example of this possibility arises within the context of theories with **anomalous $U(1)_H$ horizontal symmetries**. It has been shown [53] how one can choose the symmetry in such a way that all 45 tri-linear R-parity violating couplings are forbidden by holomorphy. The symmetry also provides a common origin for the supersymmetric μ term and the solution of the solar and atmospheric neutrino anomalies, potentially explaining also the required hierarchy between these.

Alternatively, if R parity is violated spontaneously [56] necessarily one obtains the bi-linear RPV model (BRPV) as the effective low-energy theory. Thus the BRPV model is the “common denominator” of the three most attractive routes in Fig. 11, “closed” under the renormalization group. It also constitutes the simplest and most predictive extension of the MSSM which renders a very systematic investigation of R parity violation effects not only in neutrino physics but also for supersymmetry phenomenology at collider experiments.

On general grounds the spectrum of the BRPV model is expected to be hierarchical, since only one of the three neutrinos picks up mass at the tree level, by mixing with neutralinos, as indicated in Fig. 12 (left panel). Their degeneracy is lifted only by calculable loop effects (right panel) [52] due to supersymmetric particle exchanges [57]. Both the tree level atmospheric scale as well as the more complex loop-induced solar mass scale depend quadratically on RPV parameters.



Fig. 12. Neutrino masses from RPV supersymmetry. The degeneracy of the massless states on the left is lifted by supersymmetric loop effects, illustrated in the right panel.

On the other hand neutrino mixing angles are given as RPV parameter ratios. For example the left panel in Fig. 14 shows the correlation between $\tan^2_{\theta_A}$ with the Λ_2/Λ_3 ratio. Similar correlations exist expressing the solar and reactor angles as suitable RPV ratios [52].

In summary, the BRPV model provides a powerful dynamical scheme for neutrino masses which can account for the observed atmospheric and solar neutrino anomalies in terms large angle neutrino oscillations, as indicated by current data.

6 Neutrino Mixing Tests at Colliders [58,59]

Despite the smallness of neutrino masses indicated by the oscillation interpretation of current neutrino data, R-parity violation can be observable at high-energy collider experiments such as the LHC [60]. The first and most direct manifestation of RPV is the decay of the lightest supersymmetric particle (LSP). In many supergravity scenarios, such as the BRPV model, the LSP is typically the lightest neutralino and its mass could potentially be explored at LHC and other planned colliders. Since R parity is violated the LSP decays as

$$\chi^0 \rightarrow f_1 + \bar{f}_2 + \ell$$

where f_i denotes a quark, lepton or neutrino and ℓ denotes either a charged lepton or a neutrino. In Fig. 13 we show the typical LSP decay length expected in the BRPV model. Clearly the decay length is short enough to be studied at the LHC. Moreover the decay products are mostly visible, with the main decay channel being semi-leptonic, into two b-jets plus missing transverse momentum.

Most importantly, the BRPV model provides an unambiguous test of the mixing angles involved in the neutrino anomalies at accelerators, by measuring the decay patterns of the lightest neutralino [58]. For example, in the right panel of Fig. 14 I illustrate how the ratio of semileptonic neutralino decay branching ratios to muons and tau leptons is **predicted** by the measured value of the atmospheric mixing angle indicated by atmospheric data. Similar considerations hold if the LSP is not the lightest neutralino, but some other supersymmetric particles. For example when the stop quark is the LSP, similar correlations exist between the stop decay branching ratios and the solar angle [59].




Fig. 13. LSP decay length (in cm) vs its mass (in GeV) in the BRPV model, from [60].




Fig. 14. Testing the atmospheric angle in LSP decays, from [58].

7 Life Beyond Oscillations?

Massive neutrinos typically have Non-Standard Interactions (NSI), which can be both flavour-changing (FC) and non-universal (NU) [61].

Gauge-type NSI are defined as coming from a nontrivial structure of CC and NC weak interactions. They arise in seesaw-type schemes of neutrino mass due to the rectangular (effectively non-unitary) nature of the charged current lepton mixing matrix and correspondingly non-trivial neutral current matrix (non-diagonal in mass eigenstate neutrinos) [23]. In some models these NSI may be sizable and lead to flavor and CP violation, even with strictly degenerate massless neutrinos [62]. On the other hand FC-NSI may also be induced by the exchange of spinless bosons, a situation that arises in radiative models of neutrino mass [39] as well as low-energy SUSY models with broken R parity [54,55,56]. They also arise in some supersymmetric unified models [63], where they may be calculable by the renormalization group evolution. Finally, majoron models of neutrino mass lead to pseudoscalar NSI which may induce invisible neutrino decays [26,64].

In addition to such renormalizable NSI, gauge theories of neutrino mass may also lead to NSI of higher dimension ≥ 5 , such as Majorana neutrino transition magnetic moments which lead to the Spin-Flavour Precession phenomenon [29]. As we will see shortly, this may affect solar neutrino propagation in an important

way [65], providing two solutions to the solar neutrino puzzle, one of which is resonant [66] while the other is non-resonant [67].

In what follows I take an effective model-independent description of NSI in which they are given as dimension-6 terms of the type εG_F , as illustrated in Fig. 15, where ε specifies their sub-weak strength. Such NSI may affect the




Fig. 15. Effective description of neutrino non-standard interactions.

propagation of neutrinos in a variety of situations. For example they have been considered both in context of the solar [68,69,70,71] and atmospheric [14,72] neutrino problems, as well as in connection with astrophysics [73,74,75].

In the light of such NSI one may ask, how robust are oscillations? To make a long story short, the answer is simple, atmospheric oscillations are robust, solar are not. Let me start with the atmospheric ones.

8 Flavor Changing (FC) Neutrino-Matter Interactions

8.1 Atmospheric [14]

Flavor-changing neutrino-matter interactions in the ν_μ - ν_τ channel have been shown to account for the zenith-angle-dependent deficit of atmospheric neutrinos observed in the SuperKamiokande experiment contained events [72,76], without directly invoking neither neutrino mass, nor mixing.

However such FC explanation fails to reconcile the Super-Kamiokande contained events with Super-Kamiokande and MACRO up-going muons, due to the lack of energy dependence intrinsic of NSI conversions [77]. The most recent global analysis of the atmospheric neutrino anomaly in terms of NSI shows that a pure NSI mechanism is now ruled out at 99%, while the standard $\nu_\mu \rightarrow \nu_\tau$ oscillations provide a remarkably good description of the anomaly [14]. The robustness of the atmospheric $\nu_\mu \rightarrow \nu_\tau$ oscillation hypothesis was shown to provide the most stringent and model-independent limits on flavour-changing (FC) and non-universal (NU) neutrino interactions, as illustrated in Fig. 16. These limits are model independent as they are obtained from pure neutrino-physics processes. The stability of the neutrino oscillation solution to the atmospheric neutrino anomaly against the presence of non-standard neutrino interactions illustrates the robustness of the near-maximal atmospheric mixing and massive-neutrino hypothesis.




Fig. 16. Atmospheric bounds on non-standard neutrino interactions, from [14].

8.2 Neutrino Factories-Atmospheric [78]

The proposed neutrino factories aim at probing the lepton mixing angle θ_{13} with much better sensitivity [79] than possible today by combining reactor and atmospheric data [14]. Given the LMA solution of the solar neutrino problem, it follows that there is a chance that indeed neutrino factories (NUFACT, for short) will also probe the concurrent leptonic CP violating effects (for example, through the measurement of CP asymmetries).

The potential of such a generic NUFAC in probing non-standard neutrino-matter interactions has been determined in ref. [78]. It has been found that the sensitivity to flavour-changing (FC) NSI can be substantially improved with respect to present atmospheric neutrino data, especially at energies higher than approximately 50 GeV, where the effect of the tau mass is small, as illustrated in Fig. 17. For example, a 100 GeV NUFAC can probe FC neutrino interactions at the level of few $|\epsilon| < \text{few} \times 10^{-4}$ at 99 % C.L.

8.3 Solar [71]

Let us now turn to the solar neutrino oscillation hypothesis. It provides an excellent fit to the data and a powerful way to determine neutrino mass and mixing. But, is it solid? The lack of clear hints of spectral distortion, day-night or seasonal variation in the solar neutrino data implies that, despite the increasing weight of such rate-independent observables, they do not give an unambiguous smoking-gun signal for any particular solar neutrino conversion mechanism. Here I examine two viable alternatives which do not require neutrinos to be mixed.

I will show how one can have hybrid three-neutrino interpretations of the current neutrino data in which the atmospheric data are accounted for (mainly) by oscillations, while the solar neutrino data are understood in terms of some sort of NSI, as illustrated in Fig. 18.




Fig. 17. NUFACT sensitivities to non-standard neutrino interactions, from [78].



Fig. 18. Hybrid three-neutrino interpretations of current neutrino data. The figure illustrates solutions to the solar and atmospheric neutrino anomalies in which there is only one oscillation, the atmospheric one, while the solar conversions are due to non-standard interactions (NSI).

FC-NSI solutions to the solar neutrino problem have been considered in detail in ref. [71]. Since the NSI solution is energy-independent the spectrum is undistorted, the global analysis observables are the solar neutrino rates in all experiments and the day-night measurements. It was found that the NSI description of solar data is slightly better than that of the oscillation solution and that the allowed NSI regions are determined mainly by the rate analysis [71]. By using simplified ansätze for the NSI interactions it was explicitly demonstrated that the NSI values indicated by the solar data analysis are fully acceptable also for the atmospheric data. The required parameters are indicated in Fig. 19.

Since within such a three-neutrino scheme the solar and atmospheric neutrino sectors are connected not only by the neutrino mixing angle θ_{13} but also by the flavour-changing parameters accounting for the solar data, it follows that to some extent NSI and oscillations may be confused.




Fig. 19. Parameters of NSI solution to the solar neutrino anomaly, from [71].

8.4 Neutrino Factories-Solar [80,81]

The impact of NSI on the determination of neutrino mixing parameters at a neutrino factory using the so-called “golden channels” ν_e to ν_μ and $\bar{\nu}_e$ to $\bar{\nu}_\mu$ for the measurement of θ_{13} has been given in ref. [80,81]. It was shown how a certain combination of FC interactions in neutrino source and earth matter can give exactly the same signal as oscillations arising due to θ_{13} . This implies that information on θ_{13} can only be obtained if bounds on NSI are available. Taking into account the existing bounds on FC interactions, this leads to a drastic loss in sensitivity in θ_{13} , of at least two orders of magnitude. A near detector at a neutrino factory offers the possibility to obtain stringent bounds on some NSI parameters. Such near-site detector constitutes an essential ingredient of a neutrino factory and a necessary step towards the determination of θ_{13} and subsequent study of leptonic CP violation.

Additional signatures of theories leading to FC interactions would be the existence of sizable flavour non-conservation effects, such as $\mu \rightarrow e + \gamma$, $\mu - e$ conversion in nuclei, even in the massless neutrino limit [61].

9 Solar Spin-Flavour Neutrino Precession [66,67]

The Spin-Flavour Precession (SFP) of neutrinos [29] can be resonant in matter [65] and this leads to a solution of the solar neutrino anomaly. The magnetic field profile may be fixed self-consistently by physical requirements and by demanding it to solve the solar magneto-hydrodynamics (MHD) equations [66]. The resulting ν_e survival probability is displayed in Fig. 20 and compared with the one expected in the case of neutrino oscillation. There are two solutions: resonant




Fig. 20. Best fit oscillation vs spin flavor precession ν_e survival probability, from [67].

(RSFP) and non-resonant (NRSFP). Clearly both provide good global descriptions of the solar neutrino data. In the left panel of Fig. 21 the SFP parameters




Fig. 21. “Grand Unified” solar spin-flavor-precession/oscillation plot and resulting recoil electron spectra, including the unacceptable SMA spectrum, from [67].

that account for the solar neutrino problem are presented (the plot includes the first SNO CC result, as well as the 1258-day Super-Kamiokande data). In the MHD approach the resulting scheme has only three effective parameters: Δm^2 , $\mu B_\perp$ and the neutrino mixing angle θ . The rates-only analysis for fixed $\mu B_\perp$ slightly favors SFP solutions over oscillations. In addition to the resonant solution (RSFP), there is a non-resonant solution (NRSFP) in the “dark-side” [67]. Note that in the presence of a neutrino transition magnetic moment of 10^{-11} Bohr magneton, a magnetic field of 80 KGauss eliminates all large mixing solutions other than LMA, as they would violate the upper limit on anti-neutrinos from the Sun [82]. The right panel in Fig. 21 shows that the predicted LMA, RSFP and NRSFP recoil energy spectra are in excellent agreement with the data, in contrast to that of SMA. Finally I note that it has recently been shown

that the SFP solution of the solar neutrino problem can be distinguished from the currently favored oscillation solutions at Borexino [83].

Before closing let me also add that the possibility of fast neutrino decays [64] has also been advocated could also play a sub-leading role in the solar [84] and/or atmospheric neutrino problems [85]. The data allow for neutrino stability tests, though not specially stringent.

10 In a Nutshell

It is now very hard to dispute the need for physics beyond the Standard Model in order to explain current solar and atmospheric data. In order to also account for the LSND hint [86] indicating $\bar{\nu}_\mu$ to $\bar{\nu}_e$ oscillations with a eV-mass-squared difference in the framework of quantum field theory one needs an additional light sterile neutrino ν_s [41,45,87,88]. Currently LSND is neither confirmed nor ruled out by any other experiment. A unified global analysis of current neutrino oscillation data for 4-neutrino mass schemes has been given in [89] using data from solar and atmospheric neutrino experiments, as well as information from short-baseline experiments including LSND. The summary of current 4-neutrino oscillation parameters is given in Fig. 22. One finds that the solar and atmo-




Fig. 22. “Grand-unified” neutrino oscillation plot, adapted from ref. [89], before [5,6].

spheric data reject sterile neutrinos. For example, recent SNO data rule out conversion to pure sterile neutrino states at the $\sim 5\sigma$ level. A smaller degree of sterile neutrino rejection is found for atmospheric data [89]. For this reason, the best 2+2 model model has now changed from one with the sterile state mainly in the solar sector [88] to one in which it lies mainly in the atmospheric sector [87]. However I stress that, in a truly global fit including also the LSND data

[89], sterile neutrinos can not be ruled out, since the latter disfavor the Standard Model at 99.9% CL with respect to the best 4-neutrino model. The best-fit four-neutrino model is a 3+1 scheme, since in this case one can decouple ν_s from both solar and atmospheric sectors. However, in such a global sense, even 2+2 models of the type considered in ref. [87] can not yet be ruled out, being still allowed at the 90% CL [7]. For the moment the sterile neutrino oscillation hypothesis is therefore alive, though there is a clear tension between underground and short-baseline data. We wait anxiously for results from MiniBooNE to settle the issue [90] in the near future.

I thank Martin Hirsch for comments on the manuscript. This work was supported by the European Commission grants HPRN-CT-2000-00148 and HPMT-2000-00124, by the ESF *Neutrino Astrophysics Network* and by Spanish MCyT grant PB98-0693.

References

1. S. Fukuda *et al.* [SuperKamiokande Collaboration], Phys. Rev. Lett. **86** (2001) 5651 <http://www-sk.icrr.u-tokyo.ac.jp/doc/sk/> B. T. Cleveland *et al.*, Astrophys. J. **496**, 505 (1998); R. Davis, Prog. Part. Nucl. Phys. **32**, 13 (1994); SAGE Coll., J. N. Abdurashitov *et al.*, Phys. Rev. **C60**, 055801 (1999); <http://ewiserver.npl.washington.edu/SAGE/SAGE.html> GALLEX Coll., W. Hampel *et al.*, Phys. Lett. **B447**, 127 (1999); K. Lande and V. Gavrin and E. Belloti (GNO Coll.), Talks at *Neutrino 2000*, 15–21 June 2000, Sudbury, Canada, (<http://nu2000.sno.laurentian.ca>). Q. R. Ahmad *et al.* [SNO Collaboration], Phys. Rev. Lett. **87** (2001) 071301
2. Bahcall's page <http://www.sns.ias.edu/~jnb/> and references therein.
3. M. Kachelriess, A. Strumia, R. Tomas and J. W. F. Valle, Phys. Rev. D **65**, 073016 (2002)
4. M. C. Gonzalez-Garcia, P. C. de Holanda, C. Pena-Garay and J. W. F. Valle, Nucl. Phys. B **573** (2000) 3 [arXiv:hep-ph/9906469].
5. Q. R. Ahmad *et al.* [SNO Collaboration], arXiv:nucl-ex/0204008.
6. Q. R. Ahmad *et al.* [SNO Collaboration], arXiv:nucl-ex/0204009.
7. M. Maltoni *et al.*, in preparation
8. KamLAND page <http://www.awa.tohoku.ac.jp/html/KamLAND/>
9. Y. Fukuda *et al.*, Kamiokande Coll., Phys. Lett. B **335** (1994) 237; R. Becker-Szendy *et al.*, IMB Coll., Nucl. Phys. B (Proc. Suppl.) **38** (1995) 331; W.W.M. Allison *et al.*, Soudan Coll., Phys. Lett. B **449** (1999) 137. Super-Kamiokande Coll., Y. Fukuda *et al.*, Phys. Rev. Lett. **81** (1998) 1562; C. McGrew in *Neutrino Telescopes 2001*, Venice, Italy, March 2001; T. Toshito in *Moriond 2001*, Les Arcs, France, March 2001. MACRO Coll., M. Ambrosio *et al.*, Phys. Lett. B **434** (1998) 451; M. Spurio *et al.*, hep-ex/0101019; B. Barish, Talk given at *Neutrino 2000*, 15–21 June 2000, Sudbury, Canada, <http://nu2000.sno.laurentian.ca>. A. Surdo, Talk given at *TAUP 2001*, 8–12 September 2001, Gran Sasso, Italy (<http://www.lngs.infn.it/>).
10. G. Battistoni *et al.* Nucl. Phys. Proc. Suppl. **110** (2002) 336
11. M. C. Gonzalez-Garcia, M. Maltoni, C. Pena-Garay and J. W. F. Valle, Phys. Rev. D **63** (2001) 033005 [arXiv:hep-ph/0009350].
12. M. C. Gonzalez-Garcia and M. Maltoni, arXiv:hep-ph/0202218.

13. N. Fornengo, M. C. Gonzalez-Garcia and J. W. F. Valle, Nucl. Phys. B **580** (2000) 58 [arXiv:hep-ph/0002147] and references therein.
14. N. Fornengo, M. Maltoni, R. T. Bayo and J. W. F. Valle, Phys. Rev. D **65** (2002) 013010 [arXiv:hep-ph/0108043] and references therein.
15. M. Kachelriess, R. Tomas and J. W. F. Valle, JHEP **0101** (2001) 030
16. H. Minakata, H. Nunokawa, R. Tomas and J. W. F. Valle, arXiv:hep-ph/0112160.
17. A.Y. Smirnov, D.N. Spergel and J.N. Bahcall, Phys. Rev. **D49**, 1389 (1994)
B. Jegerlehner, F. Neubig and G. Raffelt, Phys. Rev. **D54**, 1194 (1996)
18. K. Hirata *et al.* [KAMIOKANDE-II Collaboration], Phys. Rev. Lett. **58** (1987) 1490. R. M. Bionta *et al.*, Phys. Rev. Lett. **58** (1987) 1494.
19. M. Kachelriess and R. Tomas, Phys. Rev. D **64** (2001) 073002
20. S. Weinberg, “Opening Address - Neutrinos ’78” *Proceedings, West Lafayette 1978, 1-21*; R. Barbieri, J. R. Ellis and M. K. Gaillard, Phys. Lett. B **90**, 249 (1980).
21. M. Gell-Mann, P. Ramond, R. Slansky, in *Supergravity*, ed. P. van Nieuwenhuizen and D. Freedman (North Holland, 1979); T. Yanagida, in *KEK lectures*, ed. O. Sawada and A. Sugamoto (KEK, 1979)
22. R. N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44** (1980) 91.
23. J. Schechter and J. W. F. Valle, Phys. Rev. D **22** (1980) 2227.
24. J. Schechter and J. W. F. Valle, Phys. Rev. D **23** (1981) 1666.
25. Y. Chikashige, R. N. Mohapatra and R. D. Peccei, Phys. Lett. B **98** (1981) 265.
26. J. Schechter and J. W. F. Valle, Phys. Rev. D **25** (1982) 774.
27. J. Schechter and J. W. F. Valle, Phys. Rev. D **25** (1982) 2951.
M. Doi, T. Kotani and E. Takasugi, Prog. Theor. Phys. Suppl. **83** (1985) 1.
28. P. B. Pal and L. Wolfenstein, Phys. Rev. D **25** (1982) 766.
29. J. Schechter and J. W. F. Valle, Phys. Rev. D **24** (1981) 1883 [ibid **25** (1982) 283]
30. L. Wolfenstein, Phys. Rev. D **17**, 2369 (1978).
S. P. Mikheev and A. Y. Smirnov, Sov. J. Nucl. Phys. **42** (1985) 913
31. For recent reviews see S. M. Bilenky, C. Giunti and W. Grimus, Eur. Phys. J. C **1** (1998) 247 M. C. Gonzalez-Garcia and Y. Nir, arXiv:hep-ph/0202058.
32. M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49** (1973) 652.
33. J. Schechter and J. W. F. Valle, Phys. Rev. D **21** (1980) 309.
34. C. Quigg, Nucl. Instrum. Meth. A **451** (2000) 1; C. Albright *et al.*, hep-ex/0008064;
B. Autin *et al.*, CERN-SPSC-98-30; S. Geer, Phys. Rev. D **57** (1998) 6989
[Erratum-ibid. D **59** (1998) 039903]
35. For a review on $\beta\beta_{0\nu}$ experiments see A. Morales, Nucl. Phys. Proc. Suppl. **77**, 335 (1999) [arXiv:hep-ph/9809540].
36. H. V. Klapdor-Kleingrothaus *et al.*, Mod. Phys. Lett. A **16**, 2409 (2001)
37. For a critical view of the $\beta\beta_{0\nu}$ hint see C. E. Aalseth *et al.*, arXiv:hep-ex/0202018
and F. Feruglio, A. Strumia and F. Vissani, arXiv:hep-ph/0201291.
38. H. V. Klapdor-Kleingrothaus *et al.*, [GENIUS Coll.], arXiv:hep-ph/9910205.
39. T. P. Cheng and L. F. Li, Phys. Rev. D **22** (1980) 2860. A. Zee, Phys. Lett. B **93** (1980) 389 [Err-ibid. B **95** (1980) 461]. K. S. Babu, Phys. Lett. B **203** (1988) 132.
40. For recent seesaw papers see <http://131.169.91.193/cgi-bin/spiface/find/hep/www?rawcmd=t+seesaw+and+tp+neutrino+and+d+after+1997&FORMAT=WWWBRIEF&SEQUENCE=>
41. See Carlo Giunti’s page <http://www.to.infn.it/~giunti/neutrino/>
42. P. H. Chankowski *et al.*, Phys. Rev. Lett. **86** (2001) 3488 [arXiv:hep-ph/0011150].
43. A. Osipowicz *et al.* [KATRIN Collaboration], arXiv:hep-ex/0109033.
44. O. Elgaroy *et al.*, arXiv:astro-ph/0204152.
45. D. O. Caldwell and R. N. Mohapatra, Phys. Rev. D **48** (1993) 259.


46. See e.g., A. Ioannisian, J. W. F. Valle, Phys. Lett. B **332** (1994) 93; B. Bamert, C.P. Burgess, Phys. Lett. B **329** (1994) 289; D. Caldwell and R. N. Mohapatra, Phys. Rev. D **50** (1994) 3477; D. G. Lee and R. N. Mohapatra, Phys. Lett. B **329** (1994) 463; A. S. Joshipura, Z. Phys. C **64**, 31 (1994).
47. J. A. Casas, J. R. Espinosa, A. Ibarra and I. Navarro, Nucl. Phys. B **569** (2000) 82 J. Ellis and S. Lola, Phys. Lett. B **458** (1999) 310 N. Haba, Y. Matsui, N. Okamura and M. Sugiura, Prog. Theor. Phys. **103** (2000) 145; E. Ma, hep-ph/9907503.
48. L. Wolfenstein, Phys. Lett. B **107** (1981) 77.
49. J. W. F. Valle, Phys. Rev. D **27** (1983) 1672.
50. A. de Gouvea and J. W. F. Valle, Phys. Lett. B **501** (2001) 115 [hep-ph/0010299].
51. Borexino page, <http://almime.mi.infn.it/>
52. M. Hirsch et al, Phys. Rev. D **62** (2000) 113008 [arXiv:hep-ph/0004115]. and arXiv:hep-ph/0202149. J. C. Romao et al, Phys. Rev. D **61** (2000) 071703
53. J. M. Mira et al, Phys. Lett. B **492** (2000) 81 [arXiv:hep-ph/0007266].
54. C. S. Aulakh and R. N. Mohapatra, Phys. Lett. B **119**, 136 (1982). G. G. Ross and J. W. F. Valle, Phys. Lett. B **151** (1985) 375. J. R. Ellis, G. Gelmini, C. Jarlskog, G. G. Ross and J. W. F. Valle, Phys. Lett. B **150** (1985) 142. A. Santamaria and J. W. F. Valle, Phys. Lett. B **195**, 423 (1987). Phys. Rev. Lett. **60**, 397 (1988). Phys. Rev. D **39**, 1780 (1989).
55. L. J. Hall and M. Suzuki, Nucl. Phys. B **231** (1984) 419. For more recent references see <http://131.169.91.193/cgi-bin/spiface/find/hep/www?rawcmd=find+tp+r+parity+and+violation+and+neutrino+and+not+cn+>
56. A. Masiero and J. W. F. Valle, Phys. Lett. B **251** (1990) 273. J. C. Romao, C. A. Santos and J. W. F. Valle, Phys. Lett. B **288** (1992) 311. J. C. Romao and J. W. F. Valle, Nucl. Phys. B **381** (1992) 87
57. M. A. Diaz, J. C. Romao and J. W. F. Valle, Nucl. Phys. B **524** (1998) 23 [hep-ph/9706315] and references therein.
58. W. Porod, M. Hirsch, J. Romao and J. W. F. Valle, Phys. Rev. D **63** (2001) 115004
59. D. Restrepo, W. Porod and J. W. F. Valle, Phys. Rev. D **64** (2001) 055011
60. A. Bartl et al, Nucl. Phys. B **600** (2001) 39 [hep-ph/0007157].
61. For a review see, e.g. J. W. F. Valle, Prog. Part. Nucl. Phys. **26** (1991) 91.
62. R. N. Mohapatra and J. W. F. Valle, Phys. Rev. D **34** (1986) 1642. D. Wyler and L. Wolfenstein, Nucl. Phys. B **218** (1983) 205.
63. L. J. Hall, V. A. Kostelecky and S. Raby, Nucl. Phys. B **267** (1986) 415
64. J. W. F. Valle, Phys. Lett. B **131** (1983) 87; G. Gelmini, J. W. F. Valle, Phys. Lett. B **142** (1984) 181; J. W. F. Valle, Phys. Lett. B **159** (1985) 49; A. Joshipura, S. Rindani, Phys. Rev. D **46** (1992) 3000
65. E. K. Akhmedov, Phys. Lett. B **213** (1988) 64. C. S. Lim and W. J. Marciano, Phys. Rev. D **37**, 1368 (1988).
66. O. G. Miranda et al, Nucl. Phys. B **595** (2001) 360 [arXiv:hep-ph/0005259].
67. O. G. Miranda et al, Phys. Lett. B **521** (2001) 299 [arXiv:hep-ph/0108145].
68. J.W. F. Valle, Phys. Lett. **199B** (1987) 432.
69. M.M. Guzzo et al , Phys. Lett. **B271** (1991) 17.
70. S. Bergmann et al, Phys. Rev. **D62** (2000) 073001 [hep-ph/0004049].
71. M. Guzzo et al, Nucl. Phys. B **629** (2002) 479 [arXiv:hep-ph/0112310].
72. M.C. Gonzalez-Garcia *et al.*, Phys. Rev. Lett. **82** (1999) 3202, hep-ph/9809531; N. Fornengo, M. C. Gonzalez-Garcia and J. W. F. Valle, JHEP **0007** (2000) 006
73. H. Nunokawa, Y.Z. Qian, A. Rossi and J.W. F. Valle, Phys. Rev. **D54** (1996) 4356.
74. D. Grasso, H. Nunokawa and J. W. F. Valle, Phys. Rev. Lett. **81** (1998) 2412
75. G. L. Fogli, E. Lisi, A. Mirizzi and D. Montanino, arXiv:hep-ph/0202269.

76. N. Fornengo, M. C. Gonzalez-Garcia and J. W. F. Valle, JHEP **0007** (2000) 006 [arXiv:hep-ph/9906539] and references therein.
77. G. L. Fogli et al, Nucl. Phys. Proc. Suppl. **91** (2000) 167 [arXiv:hep-ph/0009269].
78. P. Huber and J. W. F. Valle, Phys. Lett. B **523** (2001) 151 [arXiv:hep-ph/0108193].
79. M. Freund, P. Huber and M. Lindner, Nucl. Phys. B **615** (2001) 331 K. Dick, M. Freund, M. Lindner and A. Romanino, Nucl. Phys. B **562** (1999) 29 A. Cervera et al, Nucl. Phys. B **579** (2000) 17 [Erratum-ibid. B **593** (2001) 731]
80. P. Huber, T. Schwetz and J. W. F. Valle, Phys. Rev. Lett. **88** (2002) 101804
81. P. Huber, T. Schwetz and J. W. F. Valle, arXiv:hep-ph/0202048, Phys. Rev. D
82. M. Aglietta *et al.*, JETP Lett. **63** (1996) 791
83. E. K. Akhmedov and J. Pulido, Phys. Lett. B **529** (2002) 193
84. J. N. Bahcall, S. T. Petcov, S. Toshev and J. W. F. Valle, Phys. Lett. B **181** (1986) 369; for a recent paper see A. Bandyopadhyay, S. Choubey and S. Goswami, Phys. Rev. D **63** (2001) 113019 and references therein.
85. V. Barger et al, Phys. Rev. Lett. **82**, 2640 (1999) and references therein.
86. A. Aguilar *et al.* [LSND Collaboration], Phys. Rev. D **64** (2001) 112007 [arXiv:hep-ex/0104049]. <http://www.neutrino.lanl.gov/LSND/>
87. J. T. Peltoniemi, D. Tommasini and J. W. F. Valle, Phys. Lett. B **298** (1993) 383. A. Ioannisian and J. W. F. Valle, Phys. Rev. D **63** (2001) 073002 [arXiv:hep-ph/9911349]. M. Hirsch and J. W. F. Valle, Phys. Lett. B **495** (2000) 121
88. J. T. Peltoniemi and J. W. F. Valle, Nucl. Phys. B **406**, 409 (1993)
89. M. Maltoni, T. Schwetz and J. W. F. Valle, Phys. Rev. D **65** (2002) 093004 [arXiv:hep-ph/0112103].
90. MiniBooNE's page <http://www-boone.fnal.gov/>



ATM-OSC

SOL-NSI





0-0



NU-CHI-MIX

LOOPS